
No.



THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE: PRESENTS SHATE COME;

Monsanto Jechnology I. I. G.

MICCERS, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLETE WITH, AND THE THE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW SCHERIFORE THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) FOR THE TERM OF TWENTY FOLLS AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY FOLLS AND THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC LIPS AND THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE SUBJECT TO SALE, OR REPRODUCING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE PROPAGATION, OR STOCKING IT FOR ANY OF THE PROPAGATION OF USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT THE PLANT VARIETY PROTECTION ACT. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

CORN, FIELD

'1000091'

In Testimonn Whereof, I have hereunto set my hand and caused the seal of the Hunt Hariety Frotestion Office to be affixed at the City of Washington, D.C. this twenty-ninth day of April, in the year two thousand and eight.

Allasti

Q2-3-

Commissioner Plant Variety Protection Office Acricultural Marketina Service Jundy . I Jenoule

Secreta Sciculture

Patent Scientist

ST-470 (02-10-2003) designed by the Plant Variety Protection Office using Word 2009

INSTRUCTIONS

GENERAL: To be effectively filed with the Plant Variety Protection Office (PVPO), ALL of the following items must be received in the PVPO: (1) Completed application form signed by the owner; (2) completed exhibits A, B, C, E; (3) for a seed reproduced variety at least 2,500 viable untreated seeds, for a hybrid variety at least 2,500 untreated seeds of each line necessary to reproduce the variety, or for tuber reproduced varieties verification that a viable (in the sense that it will reproduce an entire plant) tissue culture will be deposited and maintained in an approved public repository; (4) check drawn on a U.S. bank for \$3,652 (\$432 filling fee and \$3,220 examination fee), payable to "Treasurer of the United States" (See Section 97.6 of the Regulations and Rules of Practice.) Partial applications will be held in the PVPO for not more than 90 days, then returned to the applicant as unfilled. Mall application and other requirements to Plant Variety Protection Office, AMS, USDA, Room 401, NAL Building, 10301 Baltimore Avenue, Beltsville, MD 20705-2351. Retain one copy for your files. All items on the face of the application are self explanatory unless noted below. Corrections on the application form and exhibits must be initialed and dated. DO NOT use masking materials to make corrections. If a certificate is allowed, you will be requested to send a check payable to "Treasurer of the United States" in the amount of \$432 for issuance of the certificate. Certificates will be issued to owner, not licensee or agent.

Plant Variety Protection Office Telephone: (301) 504-5518 FAX: (301) 504-5291

Homepage: http://www.ams.usda.gov/science/pvpo/pvp.htm

ITEM

18a. Give:

- (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method;
- (2) the details of subsequent stages of selection and multiplication;
- (3) evidence of uniformity and stability; and
- (4) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified
- 18b. Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all other varieties in the same crop. If the new variety is most similar to one variety or a group of related varieties:
 - (1) identify these varieties and state all differences objectively;
 - (2) attach statistical data for characters expressed numerically and demonstrate that these are clear differences; and
 - (3) submit, if helpful, seed and plant specimens or photographs (prints) of seed and plant comparisons which clearly indicate distinctness.
- 18c. Exhibit C forms are available from the PVPO Office for most crops; specify crop kind. Fill in Exhibit C (Objective Description of Variety) form as completely as possible to describe your variety.
- 18d. Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant color, disease resistance, etc.
- 18e. Section 52(5) of the Act requires applicants to furnish a statement of the basis of the applicant's ownership. An Exhibit E form is available from the PVPO.
- 19. If "Yes" is specified (seed of this variety be sold by variety name only, as a class of certified seed), the applicant MAY NOT reverse this affirmative decision after the variety has been sold and so labeled, the decision published, or the certificate issued. However, if "No" has been specified, the applicant may change the choice. (See Regulations and Rules of Practice, Section 97.103).
- 22. See Sections 41, 42, and 43 of the Act and Section 97.5 of the regulations for eligibility requirements.
- 23. See Section 55 of the Act for instructions on claiming the benefit of an earlier filing date.
- 21. CONTINUED FROM FRONT (Please provide a statement as to the limitation and sequence of generations that may be certified.)
- 22. CONTINUED FROM FRONT (Please provide the date of first sale, disposition, transfer, or use for each country and the circumstances, if the variety (including any harvested material) or a hybrid produced from this variety has been sold, disposed of, transferred, or used in the U.S. or other countries.)

Parent of a hybrid sold in the U.S. - March 2004

23. CONTINUED FROM FRONT (Please give the country, date of filing or issuance, and assigned reference number, if the variety or any component of the variety is protected by intellectual property right (Plant Breeder's Right or Patent).)

U.S. Patent Application No. 10/820,222 - filed April 6, 2004 (1000091)

NOTES: It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownership or assignment or owner's representative during the life of the application/certificate. There is no charge for filing a change of address. The fee for filing a change of ownership or assignment or any modification of owner's name is specified in Section 97.175 of the regulations. (See Section 101 of the Act, and Sections 97.130, 97.131, 97.175(h) of the Regulations and Rules of Practice.)

To avoid conflict with other variety names in use, the applicant must check the appropriate recognized authority. For example, for agricultural and vegetable crops, contact: Seed Branch, AMS, USDA, Room 213, Building 306, Beltsville Agricultural Research Center-East, Beltsville, MD 20705. Telephone: (301) 504-8089. http://www.ams.usda.gov/lsg/seed.htm

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 3.0 hours per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual crientation, marital or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice and TOO). USDA is no equal expectively equilibring the complaint.

TDD). USDA is an equal opportunity provider and employer.

ST-470 (02-10-2003) designed by the Plant Variety Protection Office with Word 2000. Replaces former versions of ST-470, which are obsolete.

EXHIBIT A (revised)

Origin and Breeding History I000091

Corn Variety I000091 was selected for its greater plant health, greater vigor, and greater combining ability.

| | Summer 1996 | The inbred line 7526* (a proprietary Asgrow inbred) was crossed to the inbred line LH185 (a line of Holden's Foundation Seeds) in nursery rows 96KNKNH:6615 and 6624. |
|---|------------------|---|
| | Winter 1996-97 | The F1 seed was grown and self-pollinated in nursery row 962CS100 256 IBPR. |
| | Winter 1996-97 | The F2 seed was grown and self-pollinated in nursery row 97WH100:1063 IPWH. |
| - | Summer 1997 | F3 ears were grown ear-to-row and self-pollinated. One ear was selected from nursery row 97PTNP2W:199 TXPV. |
| | Summer 1998 | F4 ear was grown ear-to-row and self-pollinated. One ear was selected from nursery row 98PTNP3W:71 TXPV. |
| | Winter 1998-99 | F5 ear was grown ear-to-row and self-pollinated. Three ears were selected from nursery row 98PTS100:176 IBPR. |
| | Summer 1999 | F6 ears were grown ear-to-row and self-pollinated. Three ears were selected from nursery row W5:4209 and designated as corn variety I000091 . |
| | Winter 1999-2000 | F7 ears were grown ear-to-row and self-pollinated. Two ears from nursery row 99/77S:1194 and two ears from nursery row 99/77S:1196 were selected. |
| | Summer 2000 | F8 ears were grown ear-to-row and self-pollinated. Five ears from nursery row 913:20 and 5 ears from nursery row 913:23 were selected. |
| - | Winter 2000-2001 | F9 ears were grown ear-to-row and self-pollinated. Eleven ears from nursery row 01MXM:5650 and 9 ears from nursery row 01MXM:5651 were selected. |
| | Summer 2001 | F10 ears were grown ear-to-row and self-pollinated. Final selection was completed in nursery rows 1761-1800. This selection consisted of bulking F11 ears. |

^{* -} Inbred Line 7526 is derived from public line Oh43

Statement of Stability and Uniformity

Corn variety 1000091 was coded in 1994 and has been reproduced by self pollination for the past two years and judged to be stable. Corn variety 1000091 is uniform for all traits observed.

Statement of Variants

Corn variety 1000091 shows no variants other than what would normally be expected due to environment or that would occur for almost any character during the course of repeated sexual reproduction.

EXHIBIT B (revised)

Statement of Distinctness

Monsanto Technology L.L.C. believes that Corn Variety I000091 is most similar to corn inbred LH185, an inbred developed by Holden's Foundation Seeds.

Corn Variety I000091and LH185 differ most significantly in the following traits:

| Trait | 1000091 | LH185 |
|----------------------|--------------------------|--------------|
| Hard Endosperm Color | White | Yellow |
| | (Lighter than 2.5 Y 9/2) | (2.5 Y 8/10) |

2003

| Variety | Ear Height (cm) | | |
|---------|---------------------|--|--|
| 1000091 | 78.9 | | |
| | Std Dev = 8.1, N=10 | | |
| LH185 | 35.6 | | |
| | Std Dev = 4.3, N=10 | | |
| P_Val | 0.00 | | |
| Signif. | ** | | |

2004

| Variety | Ear Height (cm) | | | |
|---------|---------------------|--|--|--|
| 1000091 | 80.1 | | | |
| | Std Dev = 8.2, N=10 | | | |
| LH185 | 37.6 | | | |
| | Std Dev = 5.1, N=10 | | | |
| P_Val | 0.00 | | | |
| Signif. | ** | | | |

Significance levels are indicated as: + = 10%, * = 5 %, ** = 1%

Corn Variety I000091 has greater ear height and a white endosperm color while comparative corn variety LH185 has a shorter ear height and yellow endosperm color.

EXHIBIT B (revised)

Description of Experimental Design

The corn varieties I000091, LH185 and B73 were grown at the Waterman, IL observation nursery in years 2003-2004 (2002-2004 for B73). The varieties were planted in 2 row plots with 15 plants per row in each of the three years. Trait data were collected on 10 random representative plants for most traits from each 2 row plot. Data on qualitative traits are usually collected on 10 plants from each 2 row plot. For Exhibit C all data were pooled and reported as means across the years for subject variety and the standard variety with standard deviation. The varieties are randomly planted in a 4.5 acre observation nursery which is located within a larger 18 acre field. Besides the observation nursery, this field consists of a research seed increase nursery and an IP seed inventory nursery. The location of each of these individual nurseries is rotated each year to a different location within the 18 acre field. Therefore subject inbreds are not planted adjacent to comparative or standard varieties and may be located in different areas of the larger field each year, therefore being influenced by spacial differences within the field. Growing conditions within the field are not uniform as there are some slight topographical variations such as lower areas which may accumulate and retain water or higher areas which are usually drier. The field is tiled and therefore a variety maybe planted close to a tile line while a comparative variety maybe planted further away and in a low spot within the field. Temporal varieties can exist as weather conditions from year to year can vary as well as planting dates can vary from year to year based on weather conditions. Weather conditions each year can vary the maturity rate of the varieties due to either favorable or unfavorable growing conditions.

Trait variability is not observed for each variety within its own test plot-plants are usually uniform and data are collected on the "most" representative plants- variability occurs due to spacial location of the test plot for that variety from year to year and to the temporal variation of weather conditions from year to year during the 2-3 years data are collected.

Waterman Research Station Weather Data 2002-2003

| Date | Average Precip. | Ave. Monthly Temp – Max. | Ave. Monthly Temp-Min | Ave. Monthly Rel. Humid | Ave. Monthly Rel. Humid – |
|-------------|--------------------|-----------------------------|--------------------------|----------------------------|------------------------------|
| | (mm) | (F°) | (F°) | Max (%) | Min (%) |
| June 2003 | 1.7 | 76 | 54 | 90.6 | 44.3 |
| July 2003 | 3.3 | 82 | 60 | 93.6 | 53.2 |
| August 2003 | 1.3 | 84 | 61 | 9.3 | 50.5 |
| Sept 2003 | 2.1 | 74 | 51 | 92.4 | 42.9 |
| June 2004 | 3.1 | 76 | 57 | 92.8 | 50.8 |
| July 2004 | 3.2 | 79 | 59 | 94.9 | 55.6 |
| August 2004 | 3.0 | 75 | 55 | 95.8 | 55.3 |
| Sept 2004 | 0.5 | 78 | 52 | 95.0 | 43.0 |

United States Department of Agriculture, Agricultural Marketing Service Science Division, Plant Variety Protection Office National Agricultural Library Building, Room 500 Beltsville, MD 20705

OBJECTIVE DESCRIPTION OF VARIETY CORN (Zea mays L.)

| Name of Applicant(s) | | TI/ | | Antonia Maria de Maria | Danis |
|---|--|--|--------------------------|--|----------------------------|
| Name of Applicant(s) Monsanto Technology L.L.C. | | | | iety Name or Temporar 1000091 | ry Designation |
| Address (Street & No., or R.F.D. No., City, State, Zip | FOR | OFFICIAL USE | | | |
| 800 N. Lindbergh Blvd. Creve Coeur, MO U.S.A. | PVPC | Number 260500155 | | | |
| Place the appropriate number that describes the variety whole numbers by adding leading zeroes if necessary. Traits designated by a '*' are considered necessary for | Completeness shoul | d he striven for | to establish | n the spaces below. | Right justify description. |
| COLOR CHOICES (Use in conjunction with Munsell color code to describe all color choices; 01-Light Green 06-Pale Yellow 11-Pink 16-Pale P 02-Medium Green 07-Yellow 12-Light Red 17-Purple 03-Dark Green 08-Yellow-Orange 13-Cherry Red 18-Colorl 04-Very Dark Green 09-Salmon 14-Red 19-White 05-Green-Yellow 10-Pink-Orange 15-Red & White 20-White | | | | 5 and #26 in Comments 21=Buff 22=Tan 23=Brown 24=Bronze 25=Variegated (De 26=Other (Describ | escribe) |
| STANDARD INBRED CHOICES (Use the most similar (in backgryellow Dent Families: Family Members B14 CM105, A632, B64, B68 B37 B37, B76, H84 B73 N192, A679, B73, NC268 C103 Mo17, Va102, Va35, A682 Oh43 A619, MS71, H99, Va26 WF9 W64A, A554, A654, Pa91 | Yellow E Co109, ND246 Oh7, T2 W117, W W182BN White De | Dent (Unrelated): ,, 32 (153R | C13, | ns based on grow-out Sweet Corn: Iowa5125, P39, 2132 Popcorn: SG1533, 4722, HP301, Pipecorn: Mo15W, Mo16W, Mo24W | |
| 1. TYPE: (describe intermediate types in Comments sect * 8 l=Sweet 2=Dent 3=Flint 4=Flour 5=Pop 6=Orna | | n 8=Flint-Dent | Standard Inbred Name B73 | | |
| 2. REGION WHERE DEVELOPED IN THE U.S.A.: * 2 1=Northwest 2=Northcentral 3=Northeast 4=Sout 6=Southwest 7=Other | heast 5=Southcen | tral | Standard S 2 | Seed Source NCRIPS | |
| 3. MATURITY (In Region Best Adaptability; show Heat Ur section): DAYS HEAT UNITS * 0 8 1 1 4 5 4.8 From emergence to 5 * 0 8 0 1 4 3 4.5 From emergence to 5 | 0% of plants in s 0% of plants in p len shed stimum edible qual | ilk ollen ity | DAYS 7 4 7 4 | HEAT UNITS 1 6 0 8.5 1 5 5 5.0 | |
| 4. PLANT: | Standard Deviatio | n Sample Size | | Standard Deviation | Sample Size |
| * 2 0 2.2 cm Plant Height (to tassel tip) | 9.2 | 20 | 2 0 9.6 | 13.2 | 30 |
| * 0 7 9, 5 cm Ear Height (to base of top ear node) | 8.2 | 20 | 6 6.5 | 6.3 | 30 |
| 0 1 3.7 cm Length of Top Ear Internode | 1.8 | 20 | 1 4.7 | 1.7 | 30 |
| _ Average Number of Tillers | | | | | |
| * 1. 0 Average Number of Ears per Stalk | 0.1 | 20 | 1.0 | 0.0 | 30 |
| 2 Anthocyanin of Brace Roots: 1=Absent 2=E | aint 3=Moderate 4 | =Dark | 2 | | |
| Application Variety Data | Pa | ge 1 | Standard I | nbred Data | |

| Application Variety Data | Page | Page 2 | | Standard Inbred Data | | |
|--|--|-------------|-------------------------------|----------------------|-------------|--|
| 5. LEAF: | Standard Deviation | Sample Size | 5 | Standard Deviation | Sample Size | |
| * 0 0 9.8 cm Width of Ear Node Leaf | 1.5 | 20 | 7.5 | 0.9 | 30 | |
| * 0 7 1.4 cm Length of Ear Node Leaf | 3.9 | 20 | 7 4.8 | 5.7 | 30 | |
| * 5. 4 Number of leaves above top ear | 0.5 | 15 | 5.5 | 0.5 | 15 | |
| 2 7. 0 degrees Leaf Angle | 11.9 | 20 | 23.5 | 3.7 | 30 | |
| (measure from 2nd leaf above ear at a | anthesis to stalk abo | ve leaf) | 0 2 (Munse | ell code 5 GY 4/8) | | |
| * 0 2 Leaf Color (Munsell code 5 GY 4/8) | | | 5 | | | |
| 2 Leaf Sheath Pubescence(Rate on scale | | ch fuzz) | 6 | | | |
| 4 Marginal Waves (Rate on scale from 1= | | | 6 | | | |
| 4 Longitudinal Creases (Rate on scale i | from 1=none to 9=many |) | | | | |
| 6. TASSEL: | Standard Deviation | Sample Size | S | Standard Deviation | Sample Size | |
| * 8.2 Number of Primary Lateral Branches | 2.1 | 20 | 5. 5 | 0.9 | 30 | |
| 2 3. 0 Branch Angle from Central Spike | 6.4 | 20 | 2 8. 0 | 8.4 | 30 | |
| * 3 2.4 cm Tassel Length (from top leaf collar to tassel tip) | 1.9 | 20 | 4 5.6 | 2.6 | 30 | |
| 4. 4 Pollen Shed (Rate on scale from 0=male ste | erile to 9=heavy shed |) | 6.8 | | | |
| 0 7 Anther Color (Munsell code 2.5 Y 8/10) | | | 0 7 (Munsell | code 2.5 Y 8/10) | | |
| 0 2 Glume Color (Munsell code 5 GY 4/8) | | | 1 7 (Munsell | code 5 RP 5/8) | | |
| 1 Bar Glumes (Glume Bands): 1=Absent 2=Prese | ent | | 1 | | | |
| 7a. EAR (Unhusked Data): | | | | | | |
| * 0 5 Silk Color (3 days after emergence) (Munsell | code 2.5 GY 8/6) | | 0 7 (Munsell code 2.5 Y 8/10) | | | |
| 0 2 Fresh Husk Color (25 days after 50% silking) | (Munsell code 5 GY | 1/8) | 0 2 (Munsell code 5GY 4/8) | | | |
| 2 1 Dry Husk Color (65 days after 50% Silking) (| Munsell code 2.5 Y 8/ | '4) | 2 1 (Munsell | code 2.5 Y 8/4) | | |
| * 1 Position of Ear at Dry Husk Stage: 1=Upright | 2=Horizontal 3=Pende | nt | 1 | | | |
| 8 Husk Tightness (Rate on scale from 1=very lo | ose to 9=very tight) | | 9 | | | |
| 1 Husk Extension (at harvest): 1=Short (ears e 3=Long (8-10 cm beyond ear t | xposed) 2=Medium (<8 ip) 4=Very Long (>10 | cm) | 2 | | | |
| 7b. EAR (Husked Ear Data): | Standard Deviation | Sample Size | S | tandard Deviation | Sample Size | |
| * 1 2.8 cm Ear Length | 1.4 | 15 | 1 3.7 | 0.4 | 30 | |
| * 4 1.8 mm Ear Diameter at mid-point | 1.7 | 15 | 4 4.4 | 1.1 | 15 | |
| 1 2 6.3 gm Ear Weight | 5.2 | 15 | 1 2 8.7 | 6.5 | 15 | |
| * 1 4 Number of Kernel Rows | 0.9 | 15 | 1 7.6 | 1.7 | 15 | |
| 2 Kernel Rows: 1=Indistinct 2=Distinct | | | 2 | | | |
| 1 Row Alignment: 1=Straight 2=Slightly Cu | rved 3=Spiral | | 1 | | | |
| 0 9.7 cm Shank Length | 2.3 | 15 | 7.7 | 2.6 | 15 | |
| 2 Ear Taper: 1=Slight 2=Average 3=Extreme | | | 2 | | | |
| Application Variety Data | oplication Variety Data | | | | | |
| Note: Use chart on first page to choose color codes for | r color traits. | | | | | |

| Application Variety Data | | | | T. J | | |
|--|---------------------------|--------------|--|-------------------------|-------------|--|
| | Page | Page 3 | | Standard Inbred Data | | |
| 8. KERNEL (Dried): | Standard Deviation | Sample Size | | Standard Deviation | Sample Size | |
| 1 0.7 mm Kernel Length | 0.5 | 15 | 1 1.7 | 0.1 | 15 | |
| 0 8.2 mm Kernel Width | 0.2 | 15 | 7.8 | 0.3 | 15 | |
| 0 4.3 mm Kernel Thickness | 0.2 | 15 | 4.0 | 0.2 | 15 | |
| 3 0.1 % Round Kernels (Shape Grade) | 4.4 | 500g | 3 8.7 | 6.4 | 500g | |
| 1 Aleurone Color Pattern: 1=Homozygous 2 | 2=Segregating | | 1 . | | | |
| (*) 1 9 Aleurone Color (Munsell code Lighter t | than 2.5 Y 9/2) | | 1 9 (Munse | ell code Lighter than : | 2.5 Y 9/2) | |
| * 1 9 Hard Endosperm Color (Munsell code Lic | hter than 2.5 Y 9/2) | | 2 6 (oran | ge) (Munsell code 7.5 | YR 7/8) | |
| * 0 3 Endosperm Type: 1=Sweet (sul) 2=Extra 4=High Amylose Starch 5=Waxy Starch 6 8=Super Sweet (se) 9=High Oil 10=Othe | =High Protein 7=High Lv | arch sine | 0 3 | | | |
| 2 4. 6 gm Weight per 100 Kernels (unsized sam | ple) 1.2 | 1800 seeds | 2 3.1 | 0.6 | 2000 seeds | |
| 9. COB: | Standard Deviation | Sample Size | | Standard Deviation | Sample Size | |
| * 2 6.0 mm Cob Diameter at mid-point | 1.3 | 15 | 2 7.1 | 1.7 | 15 | |
| 1 9 Cob Color (Munsell code Lighter Than 5 | Y 9/1) | | 1 1 (Munse | 11 code 5 R 6/6) | | |
| 10. DISEASE RESISTANCE (Rate from 1 (most susceptib | le) to 9 (most resistant | r): | | | | |
| leave blank if not tested; leave Race or St | rain Options blank if pe | olygenic): | | | | |
| A. Leaf Blights, Wilts, and Local Infection Diseas | | | | | | |
| 7 Anthracnose Leaf Blight (Colletotrichum gramini 6 Common Rust (Puccinia sorghi) Common Smut (Ustilago maydis) 7 Eyespot (Kabatiella zeae) 6 Goss's Wilt (Clavibacter michiganense spp. nebr 5 Gray Leaf Spot (Cercospora zeae-maydis) 7 Helminthosporium Leaf Spot (Bipolaris zeicola) 6 Northern Leaf Blight (Exserohilum turcicum) Rac 7 Southern Leaf Blight (Bipolaris maydis) Race 0 Southern Rust (Puccinia polysora) 6 Stewart's Wilt (Erwinia stewartii) Other (Specify) | askense) Race 2 e 2 | | 6 5 7 7 7 Race 2 5 Race 1 3 Race 0 | | | |
| B. Systemic Diseases | | | | | | |
| 3 Corn Lethal Necrosis (MCMV and MDMV) Read Smut (Sphacelotheca reiliana) Maize Chlorotic Dwarf Virus (MCDV) Maize Chlorotic Mottle Virus (MCMV) Maize Dwarf Mosaic Virus (MDMV) Strain Sorghum Downy Mildew of Corn (Peronosclerospora Other (Specify) | sorghi) | | - - Strain | | | |
| C. Stalk Rots | | | | | | |
| _ Anthracnose Stalk Rot (Colletotrichum graminico _ Diplodia Stalk Rot (Stenocarpella maydis) _ Fusarium Stalk Rot (Fusarium moniliforme) _ Gibberella Stalk Rot (Gibberella zeae) _ Other (Specify) | | | _ _ _ _ | | | |
| D. Ear and Kernel Rots | | | | | | |
| _ Aspergillus Ear and Kernel Rot (Aspergillus fla _ Diplodia Ear Rot (Stenocarpella maydis) _ Fusarium Ear and Kernel Rot (Fusarium moniliform _ Gibberella Ear Rot (Gibberella zeae) _ Other (Specify) | ne) | | _ _ _ _ | | 11-ains | |
| Application Variety Data | | | Standard | Inbred Data | | |
| Note: Use chart on first page to choose color codes | for color traits. | | | | | |

| Application Variety Data | Pa | ge 4 | Standard Inbre | d Data | |
|---|---|--|--|------------------------------|----------------|
| 11. INSECT RESISTANCE (Rate from 1 (most susceptible) to 9 | (most resistan | nt); | | | |
| leave blank if not tested): _ Banks Grass Mite (Oligonychus pratensis) Corn Earworm (Helicoverpa zea) Leaf-Feeding | Standard Deviation | Sample Size | _ | Standard Deviation | Sample Size |
| Silk Feeding: | | | _ | | |
| Ear Damage Corn Leaf Aphid (Rhopalosiphum maidis) Corn Sap Beetle (Carpophilus dimidiatus) European Corn Borer (Ostrinia nubilalis) 1st Generation (Typically Whorl Leaf Feeding) 2nd Generation (Typically Leaf Sheath-Collar Feeding) Stalk Tunneling: |) | *************************************** | , - - - 3 5 | | |
| Fall Armyworm (Spodoptera frugiperda) Leaf-Feeding Silk-Feeding: | | | | | |
| | | | · - · · · · · · · · · · · · · · · | | |
| | | | · _ | | |
| 2. AGRONOMIC TRAITS: | | | | | |
| <pre>7 Stay Green (at 65 days after anthesis) (Rate o</pre> | on a scale from | 1=worst | 2 0 0, 0 | | |
| 0 0.0 % Pre-anthesis Brittle Snapping | | | 0 0.0 | | |
| 0 0.0 % Pre-anthesis Root Lodging | | | 0 0.0 | | |
| 0 0.0% Post-anthesis Root Lodging (at 65 days after | anthesis) | | 0 0. 0 | | |
| Kg/ha Yield of Inbred Per Se (at 12-13% grain r | | | 0 0.0 | | |
| | | | | | |
| . MOLECULAR MARKERS: (0=data unavailable; 1=data available | e but not supp | lied; 2=data su | oplied) | | |
| 0 Isozymes 0 RFLP's 0 RAPD's | | | | | |
| Butler, D.R. 1954. A System for the Classification of Corn Emerson, R.A., G.W. Beadle, and A.C. Fraser. 1935. A Summar Farr, D.F., G.F. Bills, G.P. Chamuris, A.Y. Rossman. 1989. hytopathological Society, St. Paul, MN. Inglett, G.E. (Ed.) 1970. Corn: Culture, Processing, Production, G.Gee, D.C. 1988. Maize Diseases. APS Press, St. Paul, MN. dunsell Color Chart for Plant Tissues. Macbeth. P.O. Box 23 the Mutants of Maize. 1968. Crop Science Society of America Shurtleff, M.C. 1980. Compendium of Corn Diseases. APS Pressparage, G.F., and J.W. Dudley (Editors). 1988. Corn and Companion, MI. Stringfield, G.H. Maize Inbred Lines of Ohio. Ohio A.E.S., U.S. Department of Agriculture. 1936, 1937. Yearbook. | ry of Linkage Fungi on Plan cts. Avi Publi , and Uses. Jol 150 pp. 30. Newburgh, la. Madison, WI ss, St. Paul, lorn Improvemen | Studies in Maize t and Plant Prod shing Company, W nn Wiley & Sons, N.Y. 12551-0230 MN. 105 pp. c, Third Edition | e. Cornell A.E.S., ducts in the Unite Jestport, CT. New York. | Mem. 180. d States. The A | |
| MMENTS (eg. state how heat units were calculated, standard: | | | where data was col | lected. Continu | e in Exhib: |
| at Unit Calculation: GDU = $\frac{\text{Daily Max Temp }}{2}$ (<=86°F) + $\frac{\text{Dail}}{2}$ | y Min Temp (>= | <u>50°F)</u> - 50°F | | | |
| pplemental data provided for pollen shed, ear weight, % ro ta and 2006 seed inventory data. | ound kernels ar | nd weight per 10 | 0 kernels from 20 | 06 production pa | arent test |

| REPRODUCE LOCALLY. Include form number and edition date on a | all reproductions. | FORM APPROVED - OMB No. 0581-005 |
|---|--|---|
| U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE EXHIBIT E STATEMENT OF THE BASIS OF OWNERSHIP | Application is required in order to det certificate is to be issued (7 U.S.C. 2 confidential until the certificate is issued. | termine if a plant variety protection 421). The information is held |
| 1. NAME OF APPLICANT(S) | 2. TEMPORARY DESIGNATION | 3. VARIETY NAME |
| Monsanto Technology L.L.C. | OR EXPERIMENTAL NUMBER | 1000091 |
| 4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP, and Country) | 5. TELEPHONE (Include area code) | 6. FAX (Include area code) |
| 800 N. Lindbergh Blvd. | (815) 758-9281 | (815) 758-3117 |
| Creve Couer, MO 63167 U.S.A. | 7. PVPO NUMBER | A AA E A A A E E |
| 8. Does the applicant own all rights to the variety? Mark an "X" in th | | 200500155 in. X YES NO |
| 9. Is the applicant (individual or company) a U.S. national or a U.S. b | pased company? If no, give name of co | ountry. X YES NO |
| 10. Is the applicant the original owner? | NO If no, please answer one | of the following: |
| a. If the original rights to variety were owned by individual(s), is (YES b. If the original rights to variety were owned by a company(les), YES 11. Additional explanation on ownership (Trace ownership from origin Corn Variety I000091 was originated and Technology L.L.C. By agreement between the stocky or devertion, discovery or devertibles. | is (are) the original owner(s) a U.S. base NO If no, give name of country nal breeder to current owner. Use the red developed by a breeder employer Monsanto Technology L.L.C. | ed company? Verse for extra space if needed): Ved by Monsanto and the breeder, all |
| rights to any invention, discovery or deve No rights to such invention, discovery or | elopment are assigned to Monsa development are retained by the | nto Technology L.L.C. |
| PLEASE NOTE: | | |
| Plant variety protection can only be afforded to the owners (not license | , | |
| If the rights to the variety are owned by the original breeder, that penational of a country which affords similar protection to nationals of | erson must be a U.S. national, national of the U.S. for the same genus and specie | f a UPOV member country, or s. |
| If the rights to the variety are owned by the company which employ nationals of a UPOV member country, or owned by nationals of a co- genus and species. | ed the original breeder(s), the company of the company of the original breeder (s), the origin | must be U.S. based, owned by nationals of the U.S. for the same |
| 3. If the applicant is an owner who is not the original owner, both the o | original owner and the applicant must me | et one of the above criteria. |
| The original breeder/owner may be the individual or company who dire Act for definitions. | ected the final breeding. See Section 41 | (a)(2) of the Plant Variety Protection |
| | · · · · · · · · · · · · · · · · · · · | |

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 0.1 hour per response, including the time for reviewing the instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provide and employer.